

ansanalysis

Analysis of the Autonomic Nervous System



White Paper

Scientific Quality Assurance of the ANS Analysis

Evidence Based Evaluation of the Autonomic Nervous System
Based on Short-Term Heart Rate Variability

published by Commit GmbH, Liebenburg, Germany

ANS Analysis

Scientific Quality Assurance and Standards

The entire team of Commit GmbH, from our employees, to our sales representatives and scientific consultants, strive to fulfill the utmost in quality standards and industry expectations. It is only by adhering to the highest scientific criteria and demands, what we feel is an indispensable prerequisite in the medical field, that we can ensure the continued quality of the ANS Analysis for users and potential customers.

In recent years, therapeutic analysis of the autonomic nervous system has risen exponentially in the medical community. To maintain appropriate use as an evidence based, analytical and therapeutic tool, the quality standards of data assessment, analysis, interpretation, and classification of reference values are essential.

For quality assurance of the product and to ensure the highest confidence value for the practitioner, we have published the current status of HRV-research in the following White Paper, in cooperation with our scientific consultants for heart rate variability (HRV) and the autonomic nervous system (ANS). Thereby, we would like to demonstrate our guarantee that ANS Analysis fulfills the highest possible scientific standards and requirements.

Clinical and Therapeutical Relevance

The evaluation of the autonomic nervous system (ANS) by means of (short-term) heart rate variability (HRV) represents an established clinical and analytical tool with a long history (Billman, 2011) currently having more than 1,000 related publications per year (Sassi et al., 2015). HRV assesses beat-to-beat variability of consecutive R-waves derived from ECG-measurements within a given period of time. For more than 20 years, HRV-analysis is considered an evidence based quantitative noninvasive means of assessing cardiovascular autonomic functioning (Task Force, 1996; Berntson, 1997). Established and well-defined HRV-indices draw a valid and reliable picture of ANS-functioning and enable a detailed evaluation of the impact of therapeutic treatments. Thus, distinct markers of HRV provide an important analytical tool for all medical and therapeutical fields, where an evaluation of cardiovascular autonomic functioning seems to be relevant (Kleiger et al., 2005). Most of all, HRV-analysis provides an excellent screening-tool for cardiovascular risk stratification in different clinical and therapeutic settings and populations (Thayer et al., 2007, 2010).

Methodological Background

Data Assessment and Data Analysis

The ANS Analysis assesses short-term HRV via high-resolution bi-polar chest belt detection, showing high consistency with high-resolution ECG-measurement systems in a wide range of settings and populations (e.g. Loimaala et al., 1999; Chellakumar et al., 2005; Nunan et al., 2008, 2009; Weippert et al., 2010). The bi-polar chest belt detection site used in ANS Analysis provides an RR-detection with an accuracy of ± 1 ms (Ruha et al., 1997) and satisfies guidelines for valid HRV analysis even in cases of small RR-fluctuations (Task Force, 1996; Berntson et al., 1997).

ANS Analysis calculates established and relevant HRV-indices: mean heart rate (HR), the standard deviation of all normal-to-normal RR-intervals (SDNN, measure of overall variability), the square root of the mean of the sum of the squares of differences between adjacent RR-intervals (RMSSD, measure of parasympathetic activity), the stress index as an index of the relation between height and width of RR-distribution (SI, measure of sympathetic activity) and alpha1 of detrended fluctuation analysis (measure of self-similarity and fractal organisation of ANS). These measures characterise the most important aspects of autonomic functioning in both short and long term assessments and are of prognostic relevance for different populations (e.g. Pentilla et al., 2001; Kleiger et al., 2005; Perkiomäki et al., 2005; et al.; Banzer et al., 2006, Al Haddad et al., 2011; Saboul et al., 2013; Buchheit, 2014).

With the ANS Analysis, software quality management is ensured by comparing the systems HRV indices (incl. artefact correction) with a scientific reference software (Kubios HRV 2.1, Tarvainen et al., 2014) on a regular basis. An analysis of more than 150 patients of both sexes with a wide range of clinical indications (age: 20-80 years, heart rate: 40-120 bpm) shows negligible mean differences of the main indices HR, SDNN, RMSSD and alpha1 (<0,4%) as well as a strong relation between SI and the triangular index ($r^2=0,75$) from the reference software.

Reference Values

Normal reference values in ANS Analysis are based on a systematic review and meta-analysis of short-term HRV measures in more than 20,000 healthy adults (Nunan et al., 2010). Additionally, several methodological factors contributing to large variations in HRV values and distributions are incorporated (Pikkujamsa et al., 2001; Kleiger et al., 2005; Sandercock et al., 2005; Sandercock, 2007, Buchheit, 2014). Furthermore, potential changes with age (Antelmi et al., 2004; De Meersman & Stein, 2007) as well as clinically relevant cut-off-values for risk stratification for different populations and adjustments for lifestyle factors and/or physical activity/fitness levels are considered (Kuperi et al., 1993; Liao et al., 1996, 2002; Aubert et al., 2003; Rennie et al., 2003; Schroeder et al., 2003, Buchheit et al., 2005; Thayer et al., 2007, 2010; Sandercock et al., 2008; Pivatelli et al., 2012). Therefore, the normal reference values in the ANS Analysis are applicable to a wide range of settings and subject populations.

References

- Al Haddad, H., Laursen, P. B., Chollet, D., Ahmaidi, S., & Buchheit, M. (2011). Reliability of resting and postexercise heart rate measures. *Int J Sports Med*, 32(8), 598-605.
- Antelmi, I., de Paula, R. S., Shinzato, A. R., Peres, C. A., Mansur, A. J., & Grupi, C. J. (2004). Influence of age, gender, body mass index, and functional capacity on heart rate variability in a cohort of subjects without heart disease. *Am J Cardiol*, 93(3), 381-385.
- Aubert, A. E., Beckers, F., & Ramaekers, D. (2001). Short-term heart rate variability in young athletes. *J Cardiol*, 37 Suppl 1, 85-88.
- Aubert, A. E., Seps, B. & Beckers, F. (2003). Heart rate variability in athletes. *Sports Med*, 33(12), 889-919.
- Banzer, W., Lucki, M., Bürklein, M., Rosenhagen, A., & Vogt, L. (2006). Sportmedizinische Aspekte kardialer Risikostratifizierung - Herzfrequenzvariabilität und physische Leistungsfähigkeit. *Herzschr Elektrophys*, 17(4), 197-204.
- Berntson, G. G., Bigger, J. T., Jr., Eckberg, D. L., Grossman, P., Kaufmann, P. G., Malik, M., Nagaraja, H. N., Porges, S. W., Saul, J. P., Stone, P. H. & van der Molen, M. W. (1997). Heart rate variability: origins, methods, and interpretive caveats. *Psychophysiology*, 34(6), 623-648.
- Billman, G.E. (2011). Heart rate variability - a historical perspective. *Front Physiol*. Nov 29;2:86.
- Buchheit, M. (2014). Monitoring training status with HR measures: do all roads lead to Rome? *Front Physiol*, Feb 27;5:73.
- Buchheit, M., Simon, C., Charloux, A., Doutreleau, S., Piquard, F. & Brandenberger, G. (2005). Heart rate variability and intensity of habitual physical activity in middle-aged persons. *Med Sci Sports Exerc*, 37(9), 1530-1534.
- Chellakumar, P. J., Brumfield, A., Kunderu, K., & Schopper, A. W. (2005). Heart rate variability: comparison among devices with different temporal resolutions. *Physiol Meas*, 26(6), 979-986.
- De Meersman, R.E. & Stein, P. K. (2007). Vagal modulation and aging. *Biol Psychol*, 74(2), 165-173.
- Kang, M. G., Koh, S. B., Cha, B. S., Park, J. K., Woo, J. M., & Chang, S. J. (2004). Association between job stress on heart rate variability and metabolic syndrome in shipyard male workers. *Yonsei Med J*, 45(5), 838-846.
- Kleiger, R. E., Stein, P. K., & Bigger, J. T. Jr. (2005). Heart rate variability: measurement and clinical utility. *Ann Noninvasive Electrocardiol*, 10(1), 88-101.
- Kupari, M., Virolainen, J., Koskinen, P., & Tikkanen, M. J. (1993). Short-term heart rate variability and factors modifying the risk of coronary artery disease in a population sample. *Am J Cardiol*, 72(12), 897-903.
- Liao, D., Cai, J., Barnes, R. W., Tyroler, H. A., Rautaharju, P., Holme, I., & Heiss, G. (1996). Association of cardiac autonomic function and the development of hypertension: the ARIC study. *Am J Hypertens*, 9(12Pt1), 1147-1156.
- Liao, D., Carnethon, M., Evans, G. W., Cascio, W. E., & Heiss, G. (2002). Lower heart rate variability is associated with the development of coronary heart disease in individuals with diabetes: the atherosclerosis risk in communities (ARIC) study. *Diabetes*, 51(12), 3524-3531.
- Loimaala, A., Sievanen, H., Laukkanen, R., Parkka, J., Vuori, I., & Huikuri, H. (1999). Accuracy of a novel real-time microprocessor QRS detector for heart rate variability assessment. *Clin Physiol*, 19(1), 84-88.
- Nunan, D., Jakovljevic, D. G., Donovan, G., Hodges, L. D., Sandercock, G. R. H., & Brodie, D. A. (2008). Levels of agreement for RR intervals and short-term heart rate variability obtained from the Polar S810 and an alternative system. *Eur J Appl Physiol*, 103(5), 529-537.
- Nunan, D., Donovan, G., Jakovljevic, D. G., Hodges, L. D., Sandercock, G. R. H., & Brodie, D. A. (2009). Validity and reliability of short-term heart-rate variability from the Polar S810. *Med Sci Sports Exerc*, 41(1), 243-250.
- Nunan, D., Sandercock, G. R. H., & Brodie, D. A. (2010). A quantitative systematic review of normal values for short-term heart rate variability in healthy adults. *Pacing Clin Electrophysiol*, 33(11), 1407-1417.
- Penttila, J., Helminen, A., Jartti, T., Kuusela, T., Huikuri, H. V., Tulppo, M. P., Coffeng, R. & Scheinin, H. (2001). Time domain, geometrical and frequency domain analysis of cardiac vagal outflow: effects of various respiratory patterns. *Clin. Physiol*. 21(3), 365-376.
- Perkiomaki, J. S., Makikallio, T. H., & Huikuri, H. V. (2005). Fractal and complexity measures of heart rate variability. *Clin Exp Hypertens*, 27(2-3), 149-158.
- Pikkujamsa, S. M., Makikallio, T. H., Airaksinen, K. E., & Huikuri, H. V. (2001). Determinants and interindividual variation of R-R interval dynamics in healthy middle-aged subjects. *Am J Physiol Heart Circ Physiol*, 280(3), H1400-1406.
- Pivatelli, F. C., Dos Santos, M. A., Fernandes, G. B., Gatti, M., de Abreu, L. C., Valenti, V. E., ... de Godoy, M. F. (2012). Sensitivity, specificity and predictive values of linear and nonlinear indices of heart rate variability in stable angina patients. *Int Arch Med*, 5(1), 31.
- Rennie, K. L., Hemingway, H., Kumari, M., Brunner, E., Malik, M. & Marmot, M. (2003). Effects of moderate and vigorous physical activity on heart rate variability in a British study of civil servants. *Am J Epidemiol*, 158(2), 135-143.
- Ruha, A., Sallinen, S., & Nissila, S. (1997). A real-time microprocessor QRS detector system with a 1-ms timing accuracy for the measurement of ambulatory HRV. *IEEE Trans Biomed Eng*, 44(3), 159-167.
- Saboul, D., Pialoux, V. & Hautier, C. (2013). The impact of breathing on HRV measurements: implications for the longitudinal follow-up of athletes. *Eur J Sport Sci*, 13(5):534-42.
- Sandercock, G. R. H., Bromley, P. D., & Brodie, D. A. (2005). The reliability of short-term measurements of heart rate variability. *Int J Cardiol*, 103(3), 238-247.
- Sandercock, G. R. H., Hardy-Shepherd, D., Nunan, D. & Brodie, D. (2008). The relationships between self-assessed habitual physical activity and noninvasive measures of cardiac autonomic modulation in young healthy volunteers. *J Sports Sci*, 26(11), 1171-1177.
- Sandercock, G. (2007). Normative values, reliability and sample size estimates in heart rate variability. *Clin Sci (Lond)*, 113(3), 129-130.
- Sassi, R., Cerutti, S., Lombardi, F., Malik, M., Huikuri, H. V., Peng, C. K., Schmidt, G., Yamamoto, Y. (2015). Advances in heart rate variability signal analysis: joint position statement by the e-Cardiology ESC Working Group and the European Heart Rhythm Association co-endorsed by the Asia Pacific Heart Rhythm Society. *Europace*, 17(9), 1341-1353.
- Schroeder, E. B., Liao, D., Chambless, L. E., Prineas, R. J., Evans, G. W., & Heiss, G. (2003). Hypertension, blood pressure, and heart rate variability: the Atherosclerosis Risk in Communities (ARIC) study. *Hypertension*, 42(6), 1106-1111.
- Tarvainen, M.P., Niskanen, J.P., Lipponen, J.A., Ranta-Aho, P.O. & Karjalainen, P.A. (2014). Kubios HRV-heart rate variability analysis software. *Comput Methods Programs Biomed*, 113(1), 210-220.
- Task Force of the European Society of, Pacing, and the North American Society of, & Electrophysiology. (1996). Heart rate variability. Standards of measurement, physiological interpretation, and clinical use. *Eur Heart J*, 17(3), 354-381.
- Thayer, J. F., & Lane, R. D. (2007). The role of vagal function in the risk for cardiovascular disease and mortality. *Biol Psychol*, 74(2), 224-242.
- Thayer, J. F., Yamamoto, S. S., & Brosschot, J. F. (2010). The relationship of autonomic imbalance, heart rate variability and cardiovascular disease risk factors. *Int J Cardiol*, 141(2), 122-131.
- Weippert, M., Kumar, M., Kreuzfeld, S., Arndt, D., Rieger, A., & Stoll, R. (2010). Comparison of three mobile devices for measuring R-R intervals and heart rate variability: Polar S810i, Suunto t6 and an ambulatory ECG system. *Eur J Appl Physiol*, 109(4), 779-786.

More information at:

COMMIT GmbH · Poststraße 45 · 38704 Liebenburg · Germany
Phone: +49 5346 / 912 415 · Fax: +49 5346 / 912 416
info@commitgmbh.de

COMMIT
Diagnostik- und Therapieprodukte